



PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

Linda H. MALKAS et al

Confirmation No. **5889**

Appln. No.: **10/083,576**

Group Art Unit: **0000**

Filed: **February 27, 2002**

Examiner: **Unknown**

For: **METHOD FOR PURIFYING CANCER-SPECIFIC
PROLIFERATING NUCLEAR ANTIGEN**

RECEIVED

JAN 29 2003

SUPPLEMENTAL DECLARATION OF DAVID L. MARKS

OFFICE OF PETITIONS

I the undersigned, David L. Marks, do hereby declare:

1. I, David L. Marks, am an Intellectual Property law Attorney, in TEC-COM, Office of Research and Development at the University of Maryland, Baltimore.

2. On January 9, 2003, I sent a letter to Dr. Pamela Bechtel, along with a copy of the above-captioned patent application (the specification, claims and drawings) and Declaration and Power of Attorney via FedEx Second Day Delivery, at her last known home address, 600 West Grove Parkway, Apt. 1110, Tempe, Arizona 85283-4550. The FedEx tracking number for this letter was 7912 7326 0212. A copy of this letter, and the FedEx Shipping Document therefor are attached hereto as Appendix A.

3. FedEx confirmed that on January 14, 2003, this letter was not delivered because the customer (Dr. Pamela Bechtel) was not available. A copy of the FedEx internet confirmation receipt for FedEx tracking number 7912 7326 0212 showing the same is attached hereto as Appendix B.


4. On January 22, 2003, this letter was returned to my office from FedEx. A copy of the FedEx internet confirmation receipt for FedEx tracking number 6057 5228 3419 showing the same is attached

SUPPLEMENTAL DECLARATION OF DAVID L. MARKS
U.S. Appln. No. 10/083,576

of Attorney for this application or because she can not be found or reached after diligent effort on my part to find or reach her.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.


David L. Marks


Date



UNIVERSITY OF MARYLAND

January 9, 2003

VIA FED EX 2 Day

Pamela E. Bechtel, Ph.D.
600 West Grove Parkway, Apt. 1110
Tempe, AZ 85283-4550

Re: U.S. Patent Application 10/083,576
Title: Method for Purifying Cancer-Specific
Proliferating Cell Nuclear Antigen
Our Reference: LM-2000-051

Dear Dr. Bechtel:

Enclosed please find a copy of the above referenced patent application and a Declaration/Power of Attorney for that patent application. Please sign and return the Declaration/Power of Attorney to me as soon as possible. You can fax it to me at 410-706-1066.

Thank you for your time and assistance.

Sincerely;

A handwritten signature in dark ink, appearing to read "David L. Marks", with a long horizontal flourish extending to the right.

David L. Marks


Enclosures

Malkas:LM2000-051:Patent:NP:CWPI-LM00-051-Declaration 6

RECEIVED

METHOD OF PURIFYING CANCER-SPECIFIC PROLIFERATING CELL NUCLEAR ANTIGEN

Page 1 of 3

NAME OF SOLE OR FIRST INVENTOR:			
Given Name (first and middle [if any]) Linda H.		Family Name or Surname MALKAS	
Inventor's Signature		Date	
Residence: City Indianapolis	State IN	Country USA	Citizenship United States
Mailing Address: 6256 Lakeland Blvd.			
City Indianapolis	State IN	Zip 46234	Country United States
NAME OF SECOND INVENTOR:			
Given Name (first and middle [if any]) Robert J.		Family Name or Surname HICKEY	
Inventor's Signature		Date	
Residence: City Indianapolis	State IN	Country USA	Citizenship United States
Mailing Address: 6256 Lakeland Blvd.			
City Indianapolis	State IN	Zip 46234	Country United States
NAME OF THIRD INVENTOR:			
Given Name (first and middle [if any]) Pamela E.		Family Name or Surname BECHTEL	
Inventor's Signature		Date	
Residence: City Tempe	State AZ	Country USA	Citizenship United States
Mailing Address: 600 West Grove Parkway, Apt. 110			
City Tempe	State AZ	Zip 85283	Country United States
NAME OF FOURTH INVENTOR:			
Given Name (first and middle [if any]) Min		Family Name or Surname PARK	
Inventor's Signature 		Date	
Residence: City Los Alamos	State NM	Country USA	Citizenship Korean
Mailing Address: 116 San Juan Street			
City Los Alamos	State NM	Zip 87544	Country United States
NAME OF FIFTH INVENTOR:			
Given Name (first and middle [if any]) Derek J.		Family Name or Surname HOELZ	

Mailing Address: 7755 Knickerbocker Plaza, Apt. 112

City Indianapolis State IN

Zip 46234

NAME OF SOLE OR FIRST INVENTOR:

Given Name

(first and middle [if any]) Linda H

Family Name or Surname MALKAS

Inventor's Signature

Date

Residence: City Indianapolis

State IN

Country USA

Citizenship United States

Mailing Address: 6256 Lakeland Blvd.

City Indianapolis

State IN

Zip 46234

Country United States

NAME OF SECOND INVENTOR:

Given Name

(first and middle [if any]) Robert J

Family Name or Surname HICKEY

Inventor's Signature

Date

Residence: City Indianapolis

State IN

Country USA

Citizenship United States

Mailing Address: 6256 Lakeland Blvd.

City Indianapolis

State IN

Zip 46234

Country United States

NAME OF THIRD INVENTOR:

Given Name

(first and middle [if any]) Pamela E.

Family Name or Surname BECHTEL

Inventor's Signature

Date

Residence: City Tempe

State AZ

Country USA

Citizenship United States

Mailing Address: 600 West Grove Parkway, Apt. 110

City Tempe

State AZ

Zip 85283

Country United States

NAME OF FOURTH INVENTOR:

Given Name

(first and middle [if any]) Min

Family Name or Surname PARK

Inventor's Signature

Date

Residence: City Los Alamos

State NM

Country USA

Citizenship Korean

Mailing Address: 116 San Juan Street

City Los Alamos

State NM

Zip 87544

Country United States

NAME OF FIFTH INVENTOR:

Given Name

(first and middle [if any]) Derek J

Family Name or Surname HOLLZ

City Indianapolis

State IN

Zip 46234

NAME OF SOLE OR FIRST INVENTOR:			
Given Name (first and middle [if any]) Linda H.		Family Name or Surname MALKAS	
Inventor's Signature		Date	
Residence: City Indianapolis	State IN	Country USA	Citizenship United States
Mailing Address: 6256 Lakeland Blvd.			
City Indianapolis	State IN	Zip 46234	Country United States
NAME OF SECOND INVENTOR:			
Given Name (first and middle [if any]) Robert J.		Family Name or Surname HICKEY	
Inventor's Signature		Date	
Residence: City Indianapolis	State IN	Country USA	Citizenship United States
Mailing Address: 6256 Lakeland Blvd.			
City Indianapolis	State IN	Zip 46234	Country United States
NAME OF THIRD INVENTOR:			
Given Name (first and middle [if any]) Pamela E.		Family Name or Surname BECHTEL	
Inventor's Signature		Date	
Residence: City Tempe	State AZ	Country USA	Citizenship United States
Mailing Address: 600 West Grove Parkway, Apt. 110			
City Tempe	State AZ	Zip 85283	Country United States
NAME OF FOURTH INVENTOR:			
Given Name (first and middle [if any]) Min		Family Name or Surname PARK	
Inventor's Signature		Date	
Residence: City Los Alamos	State NM	Country USA	Citizenship Korean
Mailing Address: 116 San Juan Street			
City Los Alamos	State NM	Zip 87544	Country United States
NAME OF FIFTH INVENTOR:			
Given Name (first and middle [if any]) Dong J.		Family Name or Surname HOEFL	
Inventor's Signature		Date	
Residence: City Indianapolis	State IN	Country USA	Citizenship United States
Mailing Address: 3738 Knickerbocker Place, Apt. 3D			
City Indianapolis	State IN	Zip 46234	Country United States

NAME OF SIXTH INVENTOR:			
Given Name (first and middle [if any]) Dragana		Family Name or Surname TOMIC	
Inventor's Signature		Date	
Residence: City Baltimore	State MD	Country USA	Citizenship United States
Mailing Address: 1 Fellowship Court, Apt. H			
City Baltimore	State MD	Zip 21286	Country United States
NAME OF SEVENTH INVENTOR:			
Given Name (first and middle [if any]) Lauren		Family Name or Surname SCHNAPER	
Inventor's Signature		Date	
Residence: City Lutherville	State MD	Country USA	Citizenship United States
Mailing Address: 11801 Berans Road			
City Lutherville	State MD	Zip 21093	Country United States

METHOD FOR PURIFYING CANCER-SPECIFIC
PROLIFERATING CELL NUCLEAR ANTIGEN

FIELD OF THE INVENTION

The present invention is directed to a method for purifying cancer-specific Proliferating Cell Nuclear Antigen (csPCNA), as well as to an ELISA for distinguishing csPCNA from native-proliferating cell nuclear antigen (nPCNA) and diagnosing cancer.

BACKGROUND OF THE INVENTION

10 A. Cancer

One of the least understood and most complex disease processes is the transformation that occurs as a cell becomes malignant. This process involves both genetic mutations and proteomic transformations, the result of which allows the cell to escape normal controls preventing inappropriate cell division. All cancers are unique and distinct from other cells, as well as other cancers. Despite this uniqueness, cancer cells share some common attributes. Most cancer cells proliferate outside of the normal cell cycle controls, exhibit morphological changes and exhibit various biochemical disruptions to cellular processes.

Cancer is usually diagnosed when a tumor becomes visible well after the first on-set of cellular changes. Many cancers are diagnosed after a biopsy sample is examined by histology for morphologic abnormalities, evidence of cell proliferation and genetic irregularities. There is

cellular markers for the diagnosis and

CONFIDENTIAL

prognosis of malignancies. Many of the genetic and biochemical changes occur during the early development of a tumor and these changes should be exploited for the early diagnosis of cancer.

5 Breast cancer is the leading cause of death among women in the Western world. Recent data suggests that there is a strong correlation between late detection and poor prognosis of this disease. Analysis of a thousand clinical cases indicates that
10 there is extensive genetic damage and a high rate of DNA synthesis in breast tumors in comparison with normal breast tissue. These data suggest that an alteration in the DNA replication machinery of breast cancer cells may contribute to uncontrolled
15 and error-prone DNA synthesis.

Human breast cells mediate DNA synthesis using the multiprotein replication complex termed the DNA synthesome (Coll et al, *Oncology Research*, 8:435-447 (1996)). The DNA synthesome is fully competent to
20 support in vitro DNA replication. The transformation of non-malignant human breast cells to a malignant state is accompanied by an alteration to a specific component of DNA synthesome, Proliferating Cell Nuclear Antigen (PCNA). PCNA is
25 a well-known cell-cycle marker protein, originally identified as an antigen for autoimmune disease (Bechtel et al, *Cancer Research*, 58:3264-3269 (1998)).

30 B. PCNA in Cancer/Other Cell Processes

Clinical Oncology, 121:122 (1995)). PCNA is a small (36 kD) nuclear protein involved in many cellular processes. PCNA plays crucial roles in both DNA replication and DNA repair mechanisms. PCNA has also been associated with transcription events. PCNA forms a trimer in the nucleus and acts as an accessory protein to polymerase δ , and also interacts with a variety of other proteins (Downey et al, *Cancer Cells*, 6:1211-1218 (1988)). In the evaluation of malignancy, PCNA is often used as a marker for cell proliferation. However, PCNA alone does not correlate with the stage of malignancy or the patient outcome.

A novel PCNA from breast cancer cells has been identified. The malignant breast cancer cells express a unique, acidic form of PCNA protein, i.e., csPCNA, which can clearly be distinguished from the basic form of this protein found in non-malignant cells, i.e., nPCNA. This alteration is most likely the result of a post-translational modification (Bechtel et al, *Cancer Res.*, 58:3264-3269 (1998)). However, prior to the present invention an effective method to purify csPCNA has not been described.

Recent advances in biochemical and genetic studies strongly indicate that PCNA may interact with different proteins involved in DNA mismatch repair, Okazaki fragments ligation, DNA methylation and chromatin assembly (Balajee et al, *Mutat. Res.*, 404:3-11 (1998); Ceccotti et al, *Curr Biol*: 6:1528-1531 (1996); Chen et al, *Proc Natl Acad Sci USA*, 93:1101-1106 (1996); Eki et al, *J. Biol. Chem.*, 271:14617-14622 (1996); Eki et al, *J. Biol. Chem.*, 271:14617-14622 (1996); Eki et al, *J. Biol. Chem.*, 271:14617-14622 (1996)).

Chem., 266:3087-3100 (1991); Eki et al, *J. Biol. Chem.*, 267:7284-7294 (1992); Greene et al, *Hum Mol Genet*: 8, 2263-2273 (1999); Gu et al, *Nucleic Acids Res.*, 26:1173-1178 (1998); Henderson et al, *Embo J.*,
 5 13:1450-1459 (1994); Johnson et al, *J. Biol. Chem.*, 271:27987-27990 (1996); Kelman, *Oncogene*, 14:629-640 (1997); Kolodner et al, *Curr. Opin. Genet. Dev.*, 9:89-96 (1999); Krude, *Curr. Biol.*, 9:R394-R396 (1999); Lee et al, *J. Biol. Chem.*, 266:22707-22717
 10 (1991); Levin et al, *Proc. Natl. Acad. Sci., USA*, 94:12863-12868 (1997); Levin et al, *Curr. Biol.*, 10:919-922 (2000); Martini et al, *J. Cell. Biol.*, 143:563-575 (1998); Merrill et al, *Genetics*, 148:611-624 (1998); Mimura et al, *Genes Cells*,
 15 5:439-452 (2000); Miura, *J. Radiat. Res. (Tokyo)*, 40:1-12 (1999); Moggs et al, *Mol. Cell. Biol.*, 20:1206-1218 (2000); Nishikawa et al, *Jpn. J. Cancer. Res.*, 88:1137-1142 (1997); Otterlei et al, *Embo J.*, 18:3834-3844 (1999); Pan et al, *Proc. Natl. Acad. Sci., USA*, 90:6-10 (1993); Schweitzer et al,
 20 *Genetics*, 152:953-963 (1999); Shibahara et al, *Cell*, 96:575-85 (1999); Sinicrope et al, *Clin. Cancer. Res.*, 4:1251-1261 (1998); Tom et al, *J. Biol. Chem.*, 276:24817-24825 (2001); Tomkinson et al, *Mutat. Res.*, 407:1-9 (1998); Tsurimoto, *Front. Biosci.*,
 25 4:D849-D858 (1999); Umar et al, *Cell*, 87:65-73 (1996); and Wu et al, *Nucleic Acids Res.*, 24:2036-2043 (1996)).

Xeroderma Pigmentosum (XP)G protein is reported
 30 to interact with PCNA (Gary et al, *J. Biol. Chem.*,

and the XP-G protein is reported to interact with the

PCNA-binding regions of FEN-1 and cyclin-dependent kinase inhibitor p21. XPG is a repair endonuclease similar to FEN-1 and required for nucleotide excision repair. The human XPG endonuclease cuts on
5 the 3' side of a DNA lesion, during nucleotide excision repair.

In the present invention, XPG protein was unexpectedly found to be useful in selectively purifying csPCNA, and as a part of an ELISA system
10 which can distinguish the csPCNA from the nPCNA. The detection of csPCNA in the ELISA serves as a powerful marker for early detection of malignancy.

SUMMARY OF THE INVENTION

15 An object of the present invention is to provide a method for selectively purifying csPCNA.

Another object of the present invention is to provide an ELISA for detection of csPCNA and early detection of malignancy.

20 The above-described objects, as well as others, which will be apparent from the detailed description of the invention provided hereinafter, have been met in one embodiment by a method for purifying csPCNA comprising the steps of:

25 (A) obtaining a tissue or body fluid sample comprising csPCNA;

(B) contacting said sample with a peptide comprising the amino acid sequence

30 LeuLysGlnLeuAspAlaGlnGlnThrGlnLeuArg

wherein said peptide is as defined

ORIGINAL

on a solid support and binds to said csPCNA to form a peptide-csPCNA complex; and

- 5 (C) isolating csPCNA from said peptide-csPCNA complex so as to purify said csPCNA.

In another embodiment, the above-identified objects have been met by an immunoassay for detecting csPCNA comprising:

- 10 (1) contacting a test sample with a peptide comprising the amino acid sequence LeuLysGlnLeuAspAlaGlnGlnThrGlnLeuArgIle AspSerPhePheArgLeuAlaGlnGlnGluLysGluAsp AlaLysArg (SEQ ID NO:1), which has been
15 immobilized on a solid support so as to bind csPCNA to said peptide to form a peptide-csPCNA complex; and
(2) contacting said peptide-csPCNA complex with an anti-PCNA antibody and
20 detecting binding of said antibody to said complex.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a flow chart depicting steps
25 involved in the isolation of total PCNA and csPCNA.

Figure 2 shows a Western blot of fractions collected during the isolation of PCNA. The fractions were separated on a 10% (w/v) SDS-PAGE gel, transferred to nitrocellulose and analyzed using
30 anti-PCNA antibodies. Lane 1: H₂O; Lane 2: 0.1M

Lane 9: PSW; Lane 10: PSE; Lane 11: QS;
Lane 12: XPGS; Lane 13: XPGE.

Figure 3 shows a Coomassie stained gel of fractions collected during the isolation of PCNA. The fractions were separated on a 10% (w/v) SDS-PAGE gel, and then stained with Coomassie blue. Lane 1: H; Lane 2: S1; Lane 3: S2; Lane 4: NE/S3; Lane 5: PCFT; Lane 6: PCLS; Lane 7: PCHS; Lane 8: PSFT; Lane 9: PSW; Lane 10: PSE; Lane 11: QS; Lane 12: XPGS; Lane 13: XPGE.

Figure 4 shows the results of XPGE which was subjected to 2-dimensional SDS-PAGE and Western blot analysis to identify which form of PCNA was present.

Figure 5A shows the results of a validation study on the viability of the streptavidin surface of streptavidin-coated plates.

Figure 5B shows the results of a study for determining the maximal binding capability of biotinylated XPG-GST protein to streptavidin-coated plates.

Figure 6 shows densitometric analyses of total PCNA in P4 fractions from MCF7 and MCF10A cells.

Figure 7A-7C show the results of ELISAs using MCF7 P4 and MCF10A P4 proteins.

Figure 8 shows the results of an ELISA assay comparing the abundance of csPCNA in extracts of MCF7 (malignant) cells and MCF 10A (normal) cells.

DETAILED DESCRIPTION OF THE INVENTION

As discussed above, in one embodiment of the invention, a method is provided for the detection of PCNA in a sample. The method involves the use of a PCNA antibody to detect PCNA in a sample. The method involves the use of a PCNA antibody to detect PCNA in a sample. The method involves the use of a PCNA antibody to detect PCNA in a sample.

CONFIDENTIAL

have been met by a method for purifying csPCNA comprising the steps of:

- (A) obtaining a tissue or body fluid sample comprising csPCNA;
- 5 (B) contacting said sample with a peptide comprising the amino acid sequence
LeuLysGlnLeuAspAlaGlnGlnThrGlnLeuArg
IleAspSerPhePheArgLeuAlaGlnGlnGluLys
10 GluAspAlaLysArg (SEQ ID NO:1),
wherein said peptide is immobilized on a solid support and binds to said csPCNA to form a peptide-csPCNA complex; and
- 15 (C) isolating csPCNA from said peptide-csPCNA complex so as to purify said cscNA.

Preferably, prior to step (B), the thus obtained tissue or body fluid sample of step (A) is
20 subjected to a process comprising the steps of:

- (1) homogenizing cells constituting said tissue or body fluid to obtain a homogenate (H);
- (2) separating said H into a nuclear pellet fraction (NP) and a cytosolic fraction (S1);
- 25 (3) extracting nuclei from said NP to obtain a nuclear extract (NE);
- (4) subjecting said S1 to
30 centrifugation to obtain a

(5) subjecting said S2 to centrifugation to obtain a post-mitochondrial/post-microsomal cytosolic supernatant (S3);

5 (6) combining said NE and said S3 to form an NE/S3 fraction, applying the resulting NE/S3 fraction to a phosphocellulose (a weak anion exchange matrix) column and collecting the flow
10 through (PCFT);

(7) applying the resulting PCFT to a phenylsepharose (a hydrophobic chromatography matrix) column, eluting the column with buffer comprising ethylene glycol and collecting the eluant (PSE);
15

(8) dialyzing out ethylene glycol present in the PSE to obtain a dialyzate; and

(9) applying the resulting dialyzate
20 to a Q-Sepharose (a strong anion exchange matrix) column, eluting with a dialyzate buffer comprising a salt gradient, and collecting and pooling PCNA-containing fractions to obtain said sample.

25 More preferably, the tissue or body fluid sample of step (A) further comprises native-PCNA (nPCNA), said nPCNA does not bind to said peptide in step (B) but flows through the column (XPGS), whereas csPCNA binds to said peptide in step (B) to
30 form a peptide csPCNA complex and is eluted

The particular tissue or body fluid sample which is employed in the present invention is not critical thereto. Examples of tissue which can be employed in the present invention include cervical, 5 mammary glands, esophageal, glial cells, lung, stomach, intestine, prostate, and white blood cells. Examples of body fluid which can be employed in the present invention include urine, serum and whole blood.

10 The source of the tissue or body fluid is from a subject afflicted with a cancer. The particular cancer is not critical to the present invention. The cancers can be carcinomas, sarcomas, lymphomas, or leukemias. Examples of such cancers include 15 cervical carcinoma, mammary gland carcinoma of ductal or lobular origin, gliomas, prostate, lung, esophageal, stomach, and ovarian cancer.

In step (B) the solid support employed is critical to the present invention, because the XPG 20 peptide of SEQ ID NO:1 can be expressed as a fusion protein, e.g., a GST fusion (GST = glutathione-S-transferase), and the support will depend on the fusion partner. An example of the solid support which can be employed in this case include 25 Glutathione Sepharose (Pharmacia). However, expression of the XPG-fusion protein in a Calmodulin or 6x His (oligo (6X) histidine) format can also be used in the present invention.

In step (C), csPCNA is isolated from the 30 complex by, for example, binding to a column.

through the column and appears in the flow-through

liquid exiting the column while the complex and contaminating proteins are being loaded onto the column.

The peptide represented by SEQ ID NO:1 may be synthesized chemically or by recombinant DNA techniques, as described by Gary et al, *J. Biol. Chem.*, 232:24522-24529 (1997). Further, as noted above, the peptide may be in the form of a fusion protein. The partner of the fusion protein is not critical to the present invention. Examples of such partners include Glutathione-S-Transferase (GST), Calmodulin Binding protein, and oligo(6X) histidine.

The resulting preferred csPCNA can be used to produce antibodies (monoclonal or polyclonal) specific for csPCNA by conventional techniques. The resulting purified csPCNA can also be used as standards for diagnostic kits, as well as enabling development of specific inhibitors for csPCNA (not nPCNA), the identification of the site(s) on the PCNA polypeptide that is (are) modified in nPCNA, provide the baseline for comparison to identify the type of modification sustained by nPCNA and lacking from csPCNA, the identification of specific metabolic pathways that mediate the addition or removal of this (these) post-translational modifications.

Products of antibodies specific for csPCNA can be produced by challenging mammals (e.g., rabbits, goats, horses, etc) with the peptide sequence which is post-translationally modified in nPCNA.

interdomain connective loop. They cannot distinguish

CONFIDENTIAL

csPCNA from nPCNA. Identification of the amino acid sequence of PCNA that is bound by the XPG peptide only in the csPCNA form of PCNA provides the target for preparing selective antibodies recognizing only
5 csPCNA.

Specific inhibitors of csPCNA's interaction with their target proteins can be produced by construction or expression of peptides identical to the interacting domains of csPCNA and by
10 computational chemistry methods. Through computational chemistry, the sites of interactions can be modeled and searches of existing 3-D chemical library structures or the design of new compounds can be made to disrupt and/or promote interaction
15 between csPCNA and the proteins.

Using mass spectrometry peptide analysis of tryptic fragments of csPCNA and nPCNA will identify the fragments (peptides) that are unique to csPCNA. Sequence identification of these peptides using
20 LC/MS-TDF mass spectrometry will identify the sequence, and thus the position of the modified peptides or amino acids within nPCNA that are modified in csPCNA (or that are modified in csPCNA and are not modified in nPCNA).

25 Furthermore, molecular weight and sequence analysis of these peptides by mass spectroscopy will indicate what types of post-translational modifications have been sustained by csPCNA and/or nPCNA. The identification of these
30 post-translational modifications that are unique to csPCNA

modification or the loss of the modification from

the csPCNA. In addition, computer analysis of the amino acid domains of the csPCNA that interact with the XPG peptide can reveal the presence of specific consensus sequences for a particular type of post-translational modification.

csPCNA is believed to be a better diagnostic/prognostic indicator for cancer than current methodologies which measure total PCNA. This is because total PCNA of malignant tissue/cells takes into account nPCNA and csPCNA, while this method of diagnostic/prognostic indication detects a form of PCNA only found in malignant cells/tissue. Based upon the abundance of the cancer specific form of csPCNA relative to the abundance of nPCNA the degree or extent of malignancy can be estimated. Using an ELISA assay, e.g., one whose results are shown in Figure 8, the abundance of PCNA can be measured relative to a set of standards of csPCNA of varying abundance. Comparison of the experimentally determined absorbance (using the ELISA) for the sample to the absorbance of the standards can be used to indicate the abundance of the csPCNA in the sample.

Accordingly, in another embodiment, the above-described objects of the present invention have been met by an immunoassay for detecting csPCNA comprising:

- (1) contacting a test sample with a peptide comprising the amino acid sequence

immobilized on a solid support so as to

bind csPCNA to said peptide to form a peptide-csPCNA complex; and

- 5 (2) contacting said peptide-csPCNA complex with an anti-PCNA antibody and detecting binding of said antibody to said complex.

10 The particular test sample employed is not critical to the present invention and may include any of the tissue or body fluid samples discussed above.

The particular format of the immunoassay of the present invention is not critical to the present invention. Examples of such formats include an ELISA, radio-immuno assay, dot blot assay, slot blot assay, immunoprecipitation and protein quantification, immuno-PCR, and Western blot.

15 Antibodies to PCNA can be prepared by challenging mammals (e.g., rabbits, goats, horses, etc) with the peptide sequence which is post-translationally modified in nPCNA, but not in csPCNA. All commercially available antibodies to PCNA that exist to date (~10) recognize the interdomain connector loop. They cannot distinguish csPCNA from nPCNA. Identification of the amino acid sequence of PCNA that is bound by the XPG peptide only in the csPCNA form of PCNA provides the target for preparing selective antibodies recognizing only csPCNA.

25 The detectable enzyme employed in the ELISA is

CONFIDENTIAL

As discussed above, the partner of the fusion protein is not critical to the present invention and examples of such partners include GST, Calmodulin Binding Protein and oligo(6X)histadine.

5 The particular mode of immobilization of the fusion protein on the solid support is not critical to the present invention.

It is preferable that the fusion protein is immobilized on the solid support via
10 biotin-streptavidin conjugation.

The following examples are provided for illustrative purposes only, and are in no way intended to limit the scope of the present invention.

15

EXAMPLE 1 Isolation of Total PCNA

Total PCNA was isolated using a series of centrifugation and chromatographic steps as shown in
20 Figure 1, and discussed in detail below.

A. Nuclear Extract (NE)

Human breast cancer cells, MCF7 cells (ATCC No. HTB-22), were grown in DMEM which was
25 supplemented with 10% (v/v) fetal bovine serum, 1.0% (w/w) penicillin/streptomycin, and non-essential amino acids, and then frozen until use.

13.9 g of MCF7 cells were resuspended in
30 1 volume of homogenization buffer comprising 0.5 M

5 strokes. The homogenate H was then centrifuged at 3000 rpm for 10 min, the cytosolic supernatant

removed (S1) and 10 ml of nuclear extraction buffer comprising 50 mM KCl, 50 mM HEPES (pH 7.5), 5.0 mM MgCl₂, 5.0 mM EDTA, 5.0 mM EGTA, 1.0 mM DTT and 0.1 mM PMSF, was added to the nuclear pellet.

5 The resulting nuclear pellet was rocked at 4°C for 2 hour. The nuclear pellet was centrifuged at 2,500 rpm for 10 min. The resulting supernatant was removed and centrifuged at 100,000 x g in a Ti50.2 rotor for 1 hr. The resulting supernatant, i.e.,
10 nuclear extract (NE), was collected.

EDTA and EGTA were added to S1 to 5.0 mM, and the resulting fraction was centrifuged in an SS34 rotor at 17,000 x g for 15 min. The resulting post-mitochondrial supernatant (S2) was collected,
15 and centrifuged at 100,000 x g for 1 hr in a Ti50.2 rotor. The resulting post-mitochondrial and post-microsomal cytosolic supernatant (S3), was collected.

The NE was then combined with the S3 to obtain
20 an NE/S3 fraction.

B. Phosphocellulose Column

Phosphocellulose (PC) was resuspended in a low salt buffer (LS) comprising 200 mM KCl, 50 mM HEPES
25 (pH 7.5), 5.0 mM MgCl₂, 1.0 mM DTT and 0.1 mM PMSF. A 20 ml column was poured and attached to a BioRad Biologic system. 25 ml of the NE/S3 fraction was loaded onto the column, and the flow through (PCFT) collected. A low salt fraction (PCLS) was then

... salt buffer (HS) comprising 1.0 M KCl, 50 mM HEPES

3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

(pH 7.5), 5.0 mM MgCl_2 , 1.0 mM DTT and 0.1 mM PMSF. The fractions were analyzed by Western blot analysis as described below, to determine which fractions contained PCNA.

5

C. Phenylsepharose Column

A phenylsepharose column (PS) was prepared as directed by the manufacturer (Sigma Chemical Co.) and a 6.0 ml column was poured. The column was
10 incubated with 40 ml pre-equilibration buffer comprising 20 mM potassium phosphate (pH 7.0), 0.5 mM EDTA, 0.1 mM EGTA, 10% (v/v) glycerol, 1.0 M ammonium sulfate and 1.0 mM DTT, then the identified PCNA-containing fraction from the PC column was
15 adjusted to 1.0 M ammonium sulfate and applied to the PS column. The PS column was washed with 40 ml of wash buffer comprising 20 mM potassium phosphate, 0.5 mM EDTA, 0.1 mM EGTA, 10% (v/v) glycerol, 0.5 M ammonium sulfate and 1.0 mM DTT (PSW), and the flow
20 through collected (PSFT). Fractions were eluted with 40 ml of elution buffer comprising 20 mM potassium phosphate (pH 7.0), 0.5 mM EDTA, 0.1 mM EGTA, 20% (v/v) glycerol, 10% (v/v) ethylene glycol and 1.0 mM DTT (PSE).

25 The PSE fractions were dialyzed in a 0.5 M KCl buffer comprising 0.5 M KCl, 50 mM HEPES (pH 7.5), 1.0 mM of protease inhibitor cocktail (Sigma Chemical and/or Behringer-Mannheim and/or Calbio-Chem), 5.0 mM MgCl_2 and 1.0 mM DTT; a 0.2 M
30 KCl buffer comprising 0.2 M KCl, 50 mM HEPES (pH 7.5), 1.0 mM of protease inhibitor cocktail (Sigma Chemical and/or Behringer-Mannheim and/or Calbio-Chem), 5.0 mM MgCl_2 and 1.0 mM DTT; and TDEG

Calbio-Chem, 5.0 mM MgCl_2 and 1.0 mM DTT; and TDEG

buffer comprising 50 mM Tris (pH 7.5), 1.0 mM DTT, 1.0 mM EDTA, and 10% (v/v) glycerol, and containing 100 mM KCl. The fractions were then analyzed by Western Blot, as described below, to determine which
5 fractions contained PCNA, and the PCNA-containing fractions were combined.

D. Q-Sepharose Column

A 5.0 ml Q-Sepharose column (BioRad) was
10 attached to a BioRad Biologic system. The combined PCNA-containing fractions from the PS column were concentrated to 12.5 ml, and then applied to the column. Fractions (1.0 ml) were eluted using TDEG buffer with a salt gradient of 0.1 M KCl to 0.6 M
15 KCl. The resulting fractions (QS) were analyzed by Western blot, as described below, and the PCNA-containing fractions were combined.

E. Western Blot Analysis

20 Western blot analysis was performed as described by Bechtel et al, Cancer Res., 58:3264-3269 (1998). Specifically, Western blot analysis was performed using an antibody against PCNA (Amersham) 1:1000, anti-mouse 1:3000 (Amersham)
25 and detection using chemiluminescence (Pharmacia).

The fractions collected at all of the steps of the purifications process were run on a 10% (w/v) SDS-PAGE gel, transferred to nitrocellulose and examined by Western blot analysis (Figure 2) using
30 an antibody against PCNA.

of the fractions through the phosphocellulose flow

CONFIDENTIAL

through (PCFT). Low levels of PCNA were found in phosphocellulose low salt (PCLS) and high salt (PCHS) fractions, but were undetectable in the 10 μ g/ml samples loaded onto the gel. PCNA becomes concentrated in the phenylsepharose eluant (PSE) and Q-sepharose (QS) fractions. The stained gels (Figure 3) illustrate that at each step, PCNA was further purified through the Q-sepharose column.

10

EXAMPLE 2

Isolation of csPCNA

The purified total PCNA from the Q-sepharose column was subjected to extraction using a 29 amino acid peptide (SEQ ID NO:1) derived from protein XPG, which had been fused to a GST tag. This portion of the XPG protein, i.e., amino acids 981-1009, has been shown to bind PCNA (Gary et al, *supra*).

A. Isolation of GST-XPG Fusion Protein

20 A peptide fragment of the protein XPG was cloned into *E. coli* as a GST fusion protein, as described by Gary et al, *supra*.

The transformed bacteria were grown overnight in a shaker at 37°C in 500 ml of Terrific broth comprising 23.5 g Terrific broth powder (Life Technologies Inc., Gaithersburg, MD), 2.0 ml glycerol and 100 μ g/ml Ampicillin. GST expression was induced by adding 0.1 IPTG to the culture and incubating for 4 hrs. The bacteria were collected by centrifugation in a GS lite rotor at 5000 rpm for 5 min.

The pellet was resuspended in R-PER (Pierce, Rockford, IL), rocked at room temperature for 10 min

and centrifuged at 27,000 x g for 10 min in a SS34 rotor. The supernatant, i.e., bacterial lysate, was collected.

5 Glutathione Sepharose 4B matrix beads were resuspended in the storage buffer supplied by the manufacturer, 2.0 ml of the resuspended beads were washed 10 volumes of 4°C PBS, pelleted, and then the beads were resuspended in 1.0 ml of cold PBS. The resulting GST beads were incubated with the
10 bacterial lysate for 30 min at room temperature and centrifuged in a table top centrifuge at 2500 rpm to obtain GST-XPG-glutathione beads.

B. Purification of PCNA

15 The resulting GST-XPG-glutathione beads were equilibrated in T₅₀K₃₀₀/P₁₀₀ buffer comprising 50 mM Tris (pH 7.5), 300 mM KCl and 100 mM potassium phosphate (pH 7.4). The beads were then incubated with the PCNA-containing fractions from the
20 Q-Sepharose column for 30 min at 4°C and centrifuged at 2500 rpm in a table top centrifuge. The supernatant was decanted and the beads were washed with equilibration buffer comprising 50 mM Tris (pH 7.5), 300 mM KCl and 100 mM potassium phosphate
25 (pH 7.4) (XPGS). PCNA was eluted by incubating the beads with elution buffer comprising 50 mM Tris (pH 7.5) for 30 min at 4°C, centrifuging in a microfuge and collecting the supernatant (XPGE). The wash (XPGS) and eluant (XPGE) were examined by

As shown in Figure 2, both XPGE and XPGS fractions contained PCNA. Further, as shown in Figure 3, only a single protein was present in the XPGE fraction and only a few proteins were present in the XPGS fractions.

C. 2D PAGE

The XPGE was further analyzed by 2D PAGE as described below, to determine which form of PCNA was present.

2D PAGE was performed as described by Bechtel et al, supra. Specifically, XPGE (20-40 μ g of protein) was loaded onto a first-dimension tube gel comprising 9.2 M urea, 4.0% (w/v) acrylamide, 2.0% (v/v) ampholytes (pH 3-10), and 20% (v/v) Triton X-100. The polypeptides were separated along a pH gradient created using 100 mM NaOH and 10 mM H_3PO_4 . The tube gels were then placed onto an 8.0% (w/v) acrylamide-SDS gel, and the polypeptides were resolved by molecular weight. The proteins were then examined by Western blot as described above. The results are shown in Figure 4.

As shown in Figure 4, csPNCA is the only form of PCNA present in the XPGE fraction.

D. Polymerase Assays

DNA polymerase δ activity was measured as described by Han et al, *Biochem. Pharm.*, 60:403-411 (2000). csPCNA has the ability to stimulate the

activity of DNA polymerase δ in the presence of the template, at a concentration of 0.2 OD₂₆₀ units/ml,

and the reaction mixture contained 1.0-2.0 μ g of purified DNA synthesome protein, 10 mM $MgCl_2$, 10 mM dCTP, 25 mM HEPES (pH 5.9), 200 μ g/ml of bovine serum albumin, 100 μ Ci/ml of [32 P]dGTP, and 5.0% (v/v) glycerol. The poly[dG-dC]/[dG-dC] template was boiled for 5 minutes and chilled on ice prior to use in the assay. The reaction mixture containing these components was incubated at 37°C for 15 minutes, and then spotted onto Whatman DE81 filters (Whatman International Ltd, Maidstone, England). The amount of radiolabelled nucleotide bound to the filters was quantified after washing the filters with 10 ml per filter of 0.3 M NaPPi (pH 7.5); followed by washing the filters three times with 10 ml per filter of 0.1 M NH_4 formate (pH 7.4). Afterwards, the filters were given a final wash in 95% (v/v) ethanol and then air dried, placed in scintillation vials, covered with 3 ml of scintillation fluid, and placed in a Packard TriCarb 2100TR scintillation counter (Packard Instruments Co., Meriden, CT).

The results of this assay demonstrate that addition of PCNA to purified polymerase δ increases the processivity of polymerase δ (See Bravo et al, *Nature*, 326:515-517 (1987); Downey et al, *Cancer Cells*, 6:1211-1218 (1988); and Tan et al, *Proc. Natl. Acad. Sci., USA*, 90:11014 (1986)). The addition of csPCNA to purified polymerase δ was found to increase its processivity as reported

in the following experiment. The synthesome contains bound PCNA.

EXAMPLE 3
ELISA Assay

A. Isolation of GST-XPG Fusion Protein

The PCNA binding domain of XPG (SEQ ID NO:1)
5 was ligated into a pGEX-4T-1 expression vector
(Amersham Pharmacia Biotech, Inc., Piscataway, NJ).
Recombinant protein was expressed in BL21 (DE3)
Escherichia coli using 0.8 mM IPTG. Cells were
lysed using B-Per reagents (Pierce, Rockford, IL);
10 the lysate was incubated with glutathione-agarose
beads for 2 hours at 4°C and subsequently
centrifuged for 10 minutes at 3000 rpm. The lysate
was washed twice with PBS followed by elution with
10 mM reduced glutathione (Sigma, Co., St. Louis,
15 MO) in 50 mM Tris-HCl (pH 7.4).

The resulting purified XPG-GST was biotinylated
using a commercial ECL protein biotinylation kit
(Amersham Pharmacia Biotech, Inc., Piscataway, NJ).
Briefly, 1.0 mg of protein was diluted in 1.0 ml of
20 biocarbonate buffer (pH 8.6), and incubated for
1 hour at room temperature, in 30 μ l of
biotinylation reagent per mg of protein. After
incubation, the protein sample was applied to a
Sephadex G25 column and eluted with 5.0 ml of PBS
25 (pH 7.4). Fractions of biotinylated XPG-GST protein
were then collected.

The protein profile of the biotinylated XPG-GST
protein was analyzed by 12% (w/v) SDS-PAGE and
Silver-Stain procedure (Bio-Rad silver stained plus
30 kit, Bio-Rad, Hercules, CA). To determine the

was used at a 1:1000 dilution, followed by
HRP-labeled anti-goat IgG (Santa Cruz Biotechnology,

CONFIDENTIAL

Inc., Santa Cruz, CA) at a 1:12,000 dilution in 10% (v/v) blocking buffer. The presence of biotinylated fraction of XPG-GST protein was detected using streptavidin-horse radish peroxidase
5 conjugated protein (Amersham Pharmacia Biotech, Inc., Piscataway, NJ) at dilution of 1:6,000 in PBST. Immunodetection was performed using ECL Western Blotting Detection Kit (Amersham Pharmacia Biotech, Inc., Piscataway, NJ). The biotinylated
10 XPG-GST was seen as a 32 kDa discrete protein band.

To determine a suitable concentration of biotinylated XPG-GST protein which may be enough to saturate all streptavidin-binding sites on wells of a 96-well streptavidin-coated plate (Pierce,
15 Rockford, IL), a validation of the streptavidin surface of streptavidin-coated plate was first carried out using biotinylated-HRP enzyme. Specifically, different dilutions of biotinylated enzyme HRP (biotin-HRP) (Sigma, Co., St. Louis, MO)
20 stock solution were incubated for 1 hour at room temperature, washed 5 times with PBS (pH 7.4) containing 0.1% (v/v) Tween20). Then, 100 μ l of substrate solution for HRP enzyme (Sigma Co., St. Louis, MO) was added, followed by 15 minutes of
25 incubation at 25°C. To stop the reaction, 50 μ l of stop solution (0.5 M H_2SO_4) was added. Absorbance was measured at 450 nm. Streptavidin-HRP (Sigma Co., St. Louis, MO) was used at different dilutions in 0.1% (v/v) polyoxyethylenesorbitoan Monolaurate

As shown in Figure 5A, streptavidin-coated plates were found to be completely saturated by biotin-HRP even at 1:150,000 dilution per well.

Since the streptavidin-coated plate was completely saturated by biotin-HRP at 1:150,000 dilution, the same dilution of this enzyme was used for determination of a suitable concentration of biotinylated XPG-GST protein to use. Specifically, streptavidin-coated plate was first incubated for 1 hour with different concentrations of biotinylated XPG-GST protein (10-150 $\mu\text{g/ml}$) at 25°C. After washing 5 times with PBST, biotin-HRP was added in a 1:150,000 dilution. HRP-linked enzyme was detected by adding 3,3',5,5'-tetramethylbenzidine liquid substrate system for ELISA (Sigma) of substrate-reacting solution, and absorbance was measured at 450 nm. The results are shown in Figure 5B.

As shown in Figure 5B, the optimal concentration of biotinylated XPG-GST protein that was required for saturation of all of the streptavidin-binding sites on the plate was 100 $\mu\text{g/ml}$.

25 B. ELISA

A streptavidin-coated plate was incubated with 100 $\mu\text{g/ml}$ of biotinylated XPG-GST protein for 1 hour at 25°C. Then, the wells of the plate were washed with PBST and incubated with different dilutions of protein sample.

MI) breast cell lines. The protein samples were obtained as follows.

MCF7 and MCF10A were cultured according to the protocols provided by the American Type Culture Collection (ATCC). Briefly, MCF7 cell cultures were maintained in Dulbecco's Modified Essential Media (DMEM) supplemented with 5.0% (v/v) fetal bovine serum (FBS), 100 units/ml of penicillin, 100 μ g/ml of streptomycin, and 1.0% (w/v) non-essential amino acids. MCF10A cultures were maintained in DMEM/F12 supplemented with 5.0% (v/v) Ca⁺⁺ horse serum, 10 mM HEPES, 10 μ g/ml of insulin, 20 ng/ml of epidermal growth factor (EGF), 100 ng/ml of cholera enterotoxin, 0.5 μ g/ml of hydrocortisone, 100 units/ml of penicillin, and 100 μ g/ml of streptomycin. Both cell types were grown as monolayers at 37°C, in 5.0% CO₂ atmosphere. Semiconfluent (50-75%) cell cultures were harvested and washed three times with phosphate-buffered saline (PBS), and then pelleted using low-speed centrifugation, i.e., 200 x g for 5 min at 4°C. The cell pellets were stored at -80°C until use. Next, the DNA synthesesome-enriched fraction, i.e., the P4 fraction, was isolated from the pellets of the non-malignant (MCF10A) and malignant (MCF7) breast cell lines as described by Coll et al, *Oncol. Res.*, 8(10,11):435-447 (1996), and the protein concentration in the P4 fraction was determined by a colorimetric assay.

densitometric analysis. More specifically, 5-300 μ g

of the samples were resolved in 12% (w/v) SDS-PAGE and transferred to nitrocellulose membrane. Western blot analysis was performed using a monoclonal PCNA antibody (Santa Cruz Biotechnology, Inc., Santa Cruz, CA). The values after scanning Western blot fluorograms, were expressed in Arbitrary units (Au). Discrete protein bands were quantified using BIORAD GS 710 Imaging Densitometer. The results of the densitometric analyses are shown in Figure 6, which reflects the average of three independent assays, expressed as means \pm SE.

As shown in Figure 6, the results demonstrate that the P4 fraction isolated from MCF7 cells contains twice the PCNA protein than that found in the P4 fraction isolated from MCF10A cells. Even when this result was taken in consideration as discussed below, XPG peptide was still able to distinguish csPCNA in the two cell lines.

Thereafter, dilutions of the P4 fraction from MCF7 and MCF10A cells were added to the biotinylated XPG-GST coated plates. Incubation was performed overnight at 4°C in buffer comprising 20 mM Tris-HCl (pH 7.4), 60 mM NaCl, 300 mM KCl, and 100 mM KPO₄ as described by Gary et al, supra. After washing with PBST, the wells were incubated with PCNA antibody labeled with HRP enzyme, for 1 hour at room temperature with constant agitation. The HRP-conjugated monoclonal PCNA antibody was obtained from Santa Cruz Biotechnology, Inc., Santa Cruz, CA, and was used at a dilution of 1:1000.

detected with TMB (3,3',5,5' tetramethylbenzidine)

CONFIDENTIAL

(Sigma) substrate and absorbency was read at 450 nm. The results, which are shown in Figure 7A, reflect the average of three independent assays, expressed as means \pm SE.

5 As shown in Figure 7A, XPG peptide is capable of distinguishing two forms of PCNA in the ELISA assay.

Next, serial dilutions of the P4 proteins were tested in duplicate in the ELISA and the mean value
10 of absorbency was calculated and used for comparisons. Standard curves, representing the correlation between absorbency and the abundance of the malignant and non-malignant form of PCNA, were prepared and compared to each other. The results,
15 which are shown in Figure 7B (MCF7) and Figure 7C (MCF10A), represented the mean values from three independent experiments.

As shown in Figures 7B and 7C, the XPG peptide was found to have a higher binding affinity for
20 csPCNA present in MCF7 breast cell lines.

Next, MCF10A and MCF7 cells were grown as described above. Semi-confluent (50-75%) cell cultures were harvested and washed three times with PBS, and then pelleted using low-speed
25 centrifugation, i.e., 200 x g for 5 min at 4°C. The cell pellets were stored at -80°C until use. Next, the DNA synthesize-enriched fraction, i.e., the P4 faction, was isolated from the pellets of the non-malignant (MCF10A) and malignant (MCF7) breast
30 cell lines as described in the previous section.

Cells were analyzed by ELISA. Specifically, the

P4 fraction was added to biotinylated XPG-GST coated plates. Incubation was performed overnight at 4°C in the buffer comprising 20 mM Tris-HCl (pH 7.4), 60 mM NaCl, 300 mM KCl, 100 mM KPO₄ as described above. After washing with PBST, the wells were incubated with PCNA antibody (Santa Cruz Biotechnology, Inc.), labeled with HRP enzyme, for 1 hour at room temperature with constant agitation. The PCNA antibody was used at dilution of 1:500 in 10% (v/v) blocking buffer comprising PBS (pH 7.4), 0.1% (w/v) BSA and 0.05% (v/v) Tween20. HRP enzyme was detected with TMB (3,3',5,5'-tetramethylbenzidin) (Sigma) substrate and absorbency was read at 450 nm. The amount of PCNA is indicated in arbitrary units. The results are shown in Figure 8.

As shown in Figure 8, and similar to the results in Figure 6, the P4 fraction isolated from MCF7 cells contains twice the PCNA protein than that found in the P4 fraction isolated from the MCF10A cells. Even when this result was taken in consideration (where 10 µg of the P4 fraction isolated from MCF10 cells = 1 arbitrary unit PCNA; and 5.0 µg of the P4 fraction isolated from MCF7 cells = 1 arbitrary unit PCNA), XPG peptide was still able to distinguish csPCNA in the two cell lines.

While the invention has been described in detail and with reference to specific embodiments

scope thereof.

WHAT IS CLAIMED:

Claim 1. A method for purifying cancer-specific (csPCNA) comprising the steps of:

(A) obtaining a tissue or body fluid sample comprising csPCNA;

(B) contacting said sample with a peptide comprising the amino acid sequence

LeuLysGlnLeuAspAlaGlnGlnThrGlnLeuArg
IleAspSerPhePheArgLeuAlaGlnGlnGluLys
GluAspAlaLysArg (SEQ ID NO:1),

wherein said peptide is immobilized on a solid support and binds to said csPCNA to form a peptide-csPCNA complex; and

(C) isolating csPCNA from said peptide-csPCNA complex so as to purify said csPCNA.

Claim 2. The method of Claim 1, wherein prior to step (B) the thus obtained tissue or body fluid sample of step (A) is subjected to a process comprising:

(1) homogenizing cells constituting said tissue or body fluid to obtain a homogenate (H);

(2) separating said H into a nuclear pellet fraction (NP) and a cytosolic fraction (S1);

(3) extracting nuclei from said NP to obtain a nuclear extract (NE)

post-mitochondrial cytosolic supernatant (S2);

(5) subjecting said S2 to centrifugation to obtain a post-mitochondrial/post-microsomal cytosolic supernatant (S3);

(6) combining said NE and said S3 to form an NE/S3 fraction, applying the resulting NE/S3 fraction to a weak anion exchange matrix column and collecting the flow through (PCFT);

(7) applying the resulting PCFT to a hydrophobic chromatography matrix column, eluting the column with buffer comprising ethylene glycol and collecting the eluant (PSE);

(8) dialyzing out ethylene glycol present in the PSE to obtain a dialyzate; and

(9) applying the resulting dialyzate to a strong anion exchange matrix column, eluting with a dialyzate buffer comprising a salt gradient, and collecting and pooling PCNA-containing fractions to obtain said sample.

Claim 3. The method of Claim 1 or 2, wherein said tissue or body fluid sample of step (A) further comprises native PCNA (nPCNA), said nPCNA does not bind to said peptide in step (B), whereas said

cSPNA is eluted from said cSPNA-complex.

Claim 4. An immunoassay for detecting csPCNA comprising:

- (1) contacting a test sample with a peptide comprising the amino acid sequence
LeuLysGlnLeuAspAlaGlnGlnThrGlnLeuArgIle
AspSerPhePheArgLeuAlaGlnGlnGluLysGluAsp
AlaLysArg (SEQ ID NO:1), which has been
immobilized on a solid support so as to
bind csPCNA to said peptide to form a
peptide-csPCNA complex; and
- (2) contacting said peptide csPCNA complex
with an anti-PCNA antibody and
detecting binding of said antibody to
said complex.

Claim 5. The immunoassay of Claim 4, wherein
said assay is an ELISA and said antibody is labeled
with a detectable enzyme.

Claim 6. The immunoassay of Claim 5, wherein
said enzyme is horse radish peroxidase.

Claim 7. The immunoassay of Claim 5, wherein
said peptide is a fusion protein comprising said
peptide and Glutathione-S-Transferase.

Claim 8. The immunoassay of Claim 7, wherein
said fusion protein is biotinylated and
immunobilization on said solid support via
streptavidin-coated on said solid support.

METHOD FOR PURIFYING CANCER-SPECIFIC
PROLIFERATING CELL NUCLEAR ANTIGEN

A B S T R A C T

A method for purifying cancer-specific Proliferating Cell Nuclear Antigen (csPCNA) is described, as well as an immunoassay based thereon.

FIGURE 1




FIGURE 2

Western Blot Analysis of PCNA Isolation Fractions






FIGURE 3

Analysis of csPCNA Isolation Fractions



WILEY


FIGURE 4



Downloaded from www.physiology.org/journal/ajpcell by \$\${user} on 02/11/2019. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

FIGURE 5A

tion of the viability of the streptavidin surface of the streptavidin-coated plate



- 0) blank
- 1) 1:1000 000 dilution
- 2) 1:800 000 dilution
- 3) 1: 600 000 dilution
- 4) 1:400 000 dilution
- 5) 1:200 000 dilution (2 exsp)
- 5) 1:150 000 dilution
- 7) 1:100 000 dilution (2 exsp)
- 8) 1:75000 dilution
- 9) 1:50 000 dilution (2 exsp)
- 10) 1:25 000 dilution (2 exsp)
- 11) 1:10 000 dilution (2 exsp)
- 12) 1:5000 dilution
- 13) 1:1500 dilution
- 14) 1:1000 dilution
- 15) 1:100 dilution
- 16) 1:50 dilution

CONFIDENTIAL

FIGURE 5B




FIGURE 6




FIGURE 7A

ELISA ASSAY with MCF7 P4 and MCF10 P4 proteins










FIGURE 7B



Downloaded from https://www.cambridge.org/core. University of Cambridge, on 02 Jun 2020 at 14:00:00, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/S0007122600000000

FIGURE 7C



ELISA Assay for detection of Cancer Specific Form of PCNA present in MCF-7 breast cancer cell lines



From: DAWN BROSIUS (410)706-1932
UNIVERSITY OF MARYLAND, BALT
TEC-COM/ORD
515 W LOMBARD STREET, 5TH FLOOR
BALTIMORE, MD, 21201

REVENUE BARCODE

**FedEx.****To: Pam Bechtel, Ph.D. (480)756-7478**

**600 West Grove Parkway
Apartment 1110
Tempe, AZ, 852834550**

SHIP DATE: 09JAN03
WEIGHT: 1 LBS

Ref: LM-2000-051




DELIVERY ADDRESS BARCODE(FEDEX-EDR)

FedEx ** 2DAY **

TRK # 7912 7326 0212 5581

85283-AZ-US

PHX

SA MSCA

MON

A1

Deliver by:
13JAN03

Shipping Label

[Schedule Courier](#)[Find a Dropoff Location](#)[Shipping History](#)[Shipment Complete](#)[Cancel Shipment](#)[Edit Shipment Information](#)

1. Use the "Print" feature from your browser to send this page to your laser printer.
2. Fold the printed page along the horizontal line.
3. Place label in air waybill pouch and affix it to your shipment so that the barcode portion of the label can be read and scanned.

Shipment Details

To print a copy of the shipment information for your records, please click "Shipment Details".

[Shipment Details](#)

Ship a New Package

[Ship Inside U.S.](#)[Ship Outside U.S.](#)[Ship to Same Recipient](#)

Use of this system constitutes your agreement to the service conditions in the current FedEx Service Guide, available upon request. FedEx will not be responsible for any claim in excess of \$100 per package, whether the result of loss, damage, delay, non-delivery, misdelivery, or misinformation, unless you declare a higher value, pay an additional charge, document your actual loss and file a formal claim. Limitations found in the current FedEx Service Guide apply. Your right to recover from FedEx for any loss, delay, damage, or misdelivery of the package, loss of sales, income, interest, profit, attorney's fees, costs, and other forms of damage whether direct, incidental, consequential, or special is limited to the greater of \$100 or the authorized declared value. Recovery cannot exceed actual documented loss. Maximum for items of extraordinary value is \$500, e.g. jewelry, precious metals, negotiable instruments and other items listed in our Service Guide. Written claims must be filed within strict time limits. See current FedEx Service Guide.

Track shipments

Detailed results

Tracking number 791273260212
Reference number LM-2000-051
Ship date 01/09/2003
Delivered to Receipt/Fmt desk
Delivery location TEMPE AZ
Delivery date/time 01/14/2003 16:21
Signed for by K.FISH
Service type FedEx 2Day Letter

Email your detailed tracking results

From
To
To
To
Add a message to this email

Delivered TEMPE AZ	01/14/2003 16:21	Delivered to address other than recipier
Delivered TEMPE AZ	01/14/2003 16:21	Delivered to address other than recipier
Delivery attempt TEMPE AZ	01/14/2003 16:21	Customer not available or Business clo
Delivery attempt TEMPE AZ	01/14/2003 16:18	Customer not available or Business clo
Delivery attempt TEMPE AZ	01/14/2003 16:16	Customer not available or Business clo
On FedEx vehicle for delivery TEMPE AZ	01/14/2003 08:39	
Arrived at FedEx Destination Location TEMPE AZ	01/14/2003 06:51	
Package status TEMPE AZ	01/14/2003 16:17	Package in FedEx location


[Ship](#) [Rates](#) [Signature Proof](#) [Locations](#) [Pickup](#) [International Tools](#)
[Home](#) | [About FedEx](#) | [Service Info](#) | [Careers](#) | [eBusiness Tools](#) | [Manage My Account](#) | [Customer Service](#) | [Site Index](#)

Select More Online Services



Search for



- ▶ [Track Shipments](#)
- ▶ [Alternate Reference Track](#)
- ▶ [Email Track](#)
- ▶ [FedEx InSight](#)
- ▶ [Custom Critical](#)
- ▶ [Cargo Track](#)
- ▶ [FedEx Freight](#)

Related Links

- ▶ [Signature Proof](#)
- ▶ [My FedEx](#)
- ▶ [FedEx Wireless Solutions](#)
- ▶ [FedEx Address Checker](#)

Track Shipments

Detailed Results

[Quick Help](#)

Tracking Number 605752283419
Reference Number RTS 791273260212 JS
Ship Date 01/21/2003
Delivered To Receipt/Frnt desk
Delivery Location BALTIMORE MD
Delivery Date/Time 01/22/2003 08:58
Signed For By T.GILLIAM
Service Type Express Saver Pak

Tracking Options

- [Obtain a Signature Proof of Delivery](#)
- [Email these tracking results to one or more recipients](#)
- [Track More Shipments](#)



Delivered	LINTHICUM HEIGHTS MD	01/22/2003 08:58
On FedEx Vehicle or Delivery	LINTHICUM HEIGHTS MD	01/22/2003 07:59
Arrived at FedEx Destination Location	LINTHICUM HEIGHTS MD	01/22/2003 07:37
Left FedEx Ramp	BALTIMORE MD	01/22/2003 07:18
Arrived at FedEx Ramp	BALTIMORE MD	01/22/2003 06:36
Left FedEx Sort Facility	MEMPHIS TN	01/22/2003 04:51
Left FedEx Sort Facility	MEMPHIS TN	01/22/2003 00:44
Arrived at Sort Facility	MEMPHIS TN	01/21/2003 23:58
Left FedEx Ramp	PHOENIX AZ	01/21/2003 20:50
Arrived at FedEx Ramp	PHOENIX AZ	01/21/2003 19:18
Left FedEx Origin Location	TEMPE AZ	01/21/2003 19:11
Picked up by FedEx	TEMPE AZ	01/21/2003 15:23

Email Your Detailed Tracking Results

Enter your email (optional), up to three email addresses as recipients, add your message, and click on **Send Email**.

From
 To
 To
 To

Add a message to this email.

[fedex.com Terms of Use](#) | [Contact Us](#)

This site is protected by copyright and trademark laws under U.S. and International law. Review our [privacy policy](#). All rights reserved.
© 1995-2003 FedEx